

Vol. 4 (2014) No. 6

Articles

Inhibition Effect of Mace Extract Microemulsion on Vitamin C Photooxidation in Aqueous Systems

Hasbullah Hasbullah, Sri Raharjo, Pudji Hastuti

pages: 389-393 [Full text](#) DOI:10.18517/ijaseit.4.6.442

Production Cost Assessment of Palm Empty Fruit Bunch Conversion to Bio-Oil via Fast Pyrolysis

Yoga Peryoga, Maharani Dewi Solikhah, Alfonsus Agus Raksodewanto

pages: 394-400 [Full text](#) DOI:10.18517/ijaseit.4.6.443

Growth and Yield of Chili Pepper under Different Time Application of Wedelia (*Wedelia trilobata*) and Siam Weed (*Chromolaena odorata*) Organic Fertilizers

Nanik Setyowati, Zainal Muktamar, Silma Oktiasa, Dwi W Ganefiant

pages: 401-404 [Full text](#) DOI:10.18517/ijaseit.4.6.444

Regional Planning Strategic of Irrigated Agricultural Land Conversion by Considering to The Irrigation System (Case Study : Cihea Irrigation System of Cianjur)

Endang Purnama Dewi, M. Yanuar J Purwanto, Asep Sapei

pages: 405-410 [Full text](#) DOI:10.18517/ijaseit.4.6.445

Value-added and Supporting - Inhibiting Factors for the Wet Processing of Coffee

Yuli Haryati

pages: 411-415 [Full text](#) DOI:10.18517/ijaseit.4.6.446

Individual Selection in Two Population of Segregation Based on Yield and Yield component

Nurwanita Ekasari Putri, Aries Kusumawati, Irawati Chaniago, Irfan Suliansyah

pages: 416-418 [Full text](#) DOI:10.18517/ijaseit.4.6.451

Optimization of Production Xylanase from Marine Bacterium *Bacillus safensis* P20 on Sugarcane Baggase by Submerged Fermentation

Nanik Rahmani, Nadia Ulfa Jabbar Robbani, Irma Herawati Suparto, Yopi Yopi

pages: 419-422 [Full text](#) DOI:10.18517/ijaseit.4.6.458

Quality Improvement of Cassava Flour of Local Variety of Ternate Through Fermentation Method (Application on Traditional Food of North Maluku "Sagu lempeng")

Hamidin Rasulu

pages: 423-425 [Full text](#) DOI:10.18517/ijaseit.4.6.449

Vehicular Ad Hoc Network Mobility Model

Budi Rahmadya

pages: 426-429 [Full text](#) DOI:10.18517/ijaseit.4.6.457

Selection of Arbuscular Mycorrhizal Fungi (AMF) Indigenous in Ultisol for Promoting The Production of Glimin and Aggregate Formation Processes

Amrizal Saidi, Eti Farda Husin, Azwar Rasyidin, Eddiwal Eddiwal, Ismon L

pages: 430-435 [Full text](#) DOI:10.18517/ijaseit.4.6.452

Water Potential in Petanu River Estuary and Model of Water Resources Management for Sustainable Agriculture in Gianyar Regency Bali Province

Eryani I.GST AG PT, Indayati Lanya, Santosa I GST NGR, I Nyoman Norken

pages: 436-440 [Full text](#) DOI:10.18517/ijaseit.4.6.459

A Simple Purification Method of Catechin from Gambier

IJASEIT

[About](#)

[Editorial Board](#)

[Guide for Authors](#)

[Journal Contact](#)

[Online Submission](#)

[Publication Ethics](#)

[Browse Authors](#)

[NEW Update : IJASEIT in Scopus](#)

[NEW Update : IJASEIT in Scimago JR](#)

Scimago Journal Rank

International Journal on Advanced Science,...



Scopus CiteScore

1.31 2017 CiteScore

Powered by **Scopus**

Norman Ferdinal

pages: 441-443 [Full text](#) DOI:10.18517/ijaseit.4.6.460

Phenophylogy Studies in Efforts Produced Off Season Citrus (*Citrus nobilis* var, *microcarpa*)

Sulistiawati N. P. A, Rai I. N, Santosa IGN, Astarini I. A

pages: 444-449 [Full text](#) DOI:10.18517/ijaseit.4.6.462

Involvement of Government Institutions in Agroindustrial Development: A Case of Fruit Processing Industries in East Java, Indonesia

I. B. Suryaningrat

pages: 450-454 [Full text](#) DOI:10.18517/ijaseit.4.6.448

Gambier : Indonesia Leading Commodities in The Past

Hamda Fauza

pages: 455-460 [Full text](#) DOI:10.18517/ijaseit.4.6.463

Response of Rice and Carbon Emission to Application of Ameliorant Dregs in The Peat Soil with Saturation and Unsaturation Condition

Nelvia Nelvia

pages: 461-465 [Full text](#) DOI:10.18517/ijaseit.4.6.456

The Substitution of Wheat Flour with Mixed-Cassava (*Manihot utilissima*) and Red Beans-Flour (*Phaseolus vulgaris* L.) Toward The Characteristics of Instant Noodles

Novelina Novelina, Neswati Neswati, Anggun Fitria

pages: 466-469 [Full text](#) DOI:10.18517/ijaseit.4.6.464

Analysis the Oil Palm Land which can be Converted Into Paddy Fields in Irrigated Areas of Batang Tonggar

Delvi Yanti, Eri Gas Ekaputra, Roland Asmoro

pages: 470-475 [Full text](#) DOI:10.18517/ijaseit.4.6.465




Template of Journal IJASEIT

Support Contact

WAG : +62-82171-822448

E-mail : ijaseit@insightsociety.org

The Map Authors



Keywords Cloud



Published by



Member of

International Journal on Advanced Science Engineering Information Technology

[HOME](#) [ARCHIVES](#)

Editorial Team

Editor in Chief :

Sri Atmaja Putra, (Scopus ID: 26321207400), Muhammadiyah Yogyakarta University, INDONESIA

Associate Editors :

Taufik, (Scopus ID: 23670809800), California Polytechnic State University, USA
 Moustafa M Eissa, (Scopus ID: 35581693900), Helwan University, EGYPT
 Wan Mohhtar Wan Yusoff, (Scopus ID: 15019967700), Univ. Kebangsaan Malaysia, MALAYSIA
 Halimah Badioze Zaman, (Scopus ID: 25825801600), Univ. Kebangsaan Malaysia, MALAYSIA
 Son Radu, (Scopus ID: 7005251005), Universiti Putra Malaysia, Malaysia, MALAYSIA
 Mohd Razi Ismail, (Scopus ID: 25957691400), Universiti Putra Malaysia, MALAYSIA
 Takashi Oku, (Scopus ID: 56275094900), Prefectural University of Hiroshima, JAPAN
 Kohei Nakano, (Scopus ID: 7402011766), Gifu University, JAPAN
 Nurul Huda, (Scopus ID: 6701695514), Universiti Sultan Zainal Abidin, MALAYSIA
 Yandra Arkeman, (Scopus ID: 55946558300), Bogor Agriculture University, INDONESIA
 Sate Sampattagul, (Scopus ID: 7801640881), Chiangmai University, THAILAND
 Peeyush Soni, (Scopus ID: 9248907800), Asian Institute of Technology, THAILAND
 Yolanda Lechon Perez, (Scopus ID: 6502626000), Ciemat, Madrid, SPAIN
 Gabriele Arcidiacono (Scopus ID: 56656284600), G. Marconi University, ITALY
 Alessandra Pieroni (Scopus ID: 25928524500), Marconi International University, Florida - USA
 Nguyen Hay, (Scopus ID: 15834645900) Nong Lam University, VIETNAM
 Rita Muhammad Awang, (Scopus ID: 55957782400), Universiti Putra Malaysia, MALAYSIA
 Anton S Prabuwo, (Scopus ID: 18134309800), King Abdulaziz Univ, SAUDI ARABIA
 P Mangala C S De Silva, (Scopus ID: 7006461145), University of Ruhuna, SRI LANKA
 Bich Huiy Nguyen, (Scopus ID: 38191086100), Nong Lam University, VIETNAM
 Paul Kristiansen, (Scopus ID: 23097563600), University of New England, AUSTRALIA
 Amitava Basu, (Scopus ID: 21833738300), Bidhan Chandra Krishi Vidyalaya, INDIA
 Haitham Alali, (Scopus ID: 49963007000), Amman Arab University, JORDAN
 Shahrul Azman Mohd Noah, (Scopus ID: 35087633200), Universiti Kebangsaan Malaysia, MALAYSIA
 Luca Di Nunzio (Scopus ID: 57195199010), University of Rome Tor Vergata, ITALY

Managing Editor :

Rahmat Hidayat, (Scopus ID: 57191265401) Politeknik Negeri Padang, INDONESIA

Editors :

Nurhamidah, (Scopus ID: 57191636504), Andalas University, INDONESIA
 Ario Betha Juansiffero, (Scopus ID: 57189369470), Kobe University, JAPAN
 Ifa Puspasari, (Scopus ID: 55180322800), National University of Malaysia - MALAYSIA
 Zairi Ismael Rizman, (Scopus ID: 36959761800), Universiti Teknologi MARA (UiTM) (Tarengganu) MALAYSIA
 Shahreen Kasim, (Scopus ID: 36155431900), Universiti Tun Hussein Onn - MALAYSIA
 Chi-Hua Chen, (Scopus ID: 35799698800), National Chiao Tung University, TAIWAN
 Gusri Yaldi, (Scopus ID: 55359203100), Polytechnics State of Padang, INDONESIA
 Abrar Ismardi, (Scopus ID: 26633102900), Telkom University - INDONESIA

IJASEIT

[About](#)

[Editorial Board](#)

[Guide for Authors](#)

[Journal Contact](#)

[Online Submission](#)

[Publication Ethics](#)

[Browse Authors](#)

[NEW Update : IJASEIT in Scopus](#)

[NEW Update : IJASEIT in Scimago JR](#)

Scimago Journal Rank

International Journal on
Advanced Science...



Scopus CiteScore



Response of Rice and Carbon Emission to Application of Ameliorant Dregs in The Peat Soil with Saturation and Unsaturation Condition

Nevia

Department of Agrotechnology, faculty of Agriculture, Riau University, Indonesia
E-mail: nnelvia@yahoo.co.id

Abstract— Fertility of peat land is very poor and makes it not suitable for crop without any high input. This condition was indicated by the very high soil acidity (low pH), low availability of macro (N, K, Ca and P), and micro (Cu, Zn, Mn and Bo) nutrients and high cation exchange capacity (CEC) but low base saturation (BS), the presence of toxic organic acid. The main organic acids, as a result of lignin biodegradation and the sources of C-release, are of aromatic group consisting mainly of derivate phenolic acids. The peat soil material was taken at depths from 0 to 30 cm, with weathering rate saprik and dregs from Indah Kiat Pulp and Paper at Perawang, Riau. These experiments using split plot design, the main plot is the condition of the water (unsaturated and saturated) and the subplot is ameliorant dregs consisting of 4 levels (0, 10, 15 and 20 ton ha⁻¹), each combination was repeated 4 times. The activities were focused on the interaction of water condition and ameliorant dregs, and its influences to growth and yield of rice, C-emission (CO₂ and CH₄). The results showed that the application of dregs improves plant growth and increase the yield of rice (weight of dry milled grain) compared without dreg both at unsaturated condition and saturated conditions. The Carbon-release in the forms of CO₂ and CH₄ fluxes in saturated conditions is smaller than unsaturated conditions. The application dreg 10 t ha⁻¹ increase the number of productive tillers and the weight of milled rice about 35 and 75% compared without dreg on saturated condition, whereas the increase is greater unsaturated conditions about 63 and 80%. The application of dregs 10 t ha⁻¹ in saturated condition can reduce CO₂ and CH₄ emissions about 18.19% and 93.71% compared without dreg, otherwise the application dreg 10-20 t ha⁻¹ increase the production of CO₂ and CH₄ in unsaturated conditions.

Keywords— PutRice, peat soil, ameliorant dregs, CH₄ and CO₂ emission, saturated, unsaturated condition

I. INTRODUCTION

This document is a template. An electronic copy can be downloaded from the conference website. For questions on paper guidelines, please contact the conference publications committee as indicated on the conference website. Information about final paper submission is available from the conference website.

Indonesia has 188 milion ha land, including peatland about 20.9 million ha, (Wahyunto et al., 2005). Peatland fertility is very poor and makes it not suitable for crop without any high input. This condition was indicated by the very high soil acidity (low pH), low availability of macro (N, K, Ca, and P), and micro (Cu, Zn, Mn and Bo) nutrients and high cation exchange capacity (CEC) but low base saturation (BS), and the presence of toxic organic acid (Simbolon, 2009).

The main organic acid, as a result of lignin biodegradation and the sources of C-release, are of aromatic group consisting mainly of derivate phenolic acids. The concentration of such organic acids ranging from the highest

to lowest is as follows: ferulic acid \approx synapic acid > p-coumaric acid > p-hydroxybenzoic acid > vanilic acid > syringic acid (Sabiham, 2010). Phenolic acids are more phytotoxic for plants and causes stunted plant growth (Tsutuski, 1984; Stevens, *et al.*, 1994; Dohong and Sabiham, 2001), influence the biochemical and physiological processes of plants and nutrients uptake by plant (Driessen, 1978).

The drying and wetting processes on the peat materials affected the stability of organic acids, wich was indicated by loss of C- through CO₂ and CH₄ releases. The release of CO₂ and CH₄ from fibric peat was higher than that from hemic and sapric peats (Sabiham, 2010). Yagi and Minami (1990) reported that the highest rate of CH₄ emission during cultivation period (44.8 g CH₄ m⁻²) was in rice field consisting peat.

The drainage of peat release oxigen (O₂) into the surface, with promotes decomposition. Emission estimates, for land use systems with a depth of 60 cm drainage is around 55 Mg CO₂ ha⁻¹ year⁻¹ (Hooijer *et al.* 2010), based on a linear relationship between depth of water table and emissions.

Phenolic acids and C-release could be reduced to the granting of polyvalent cations such as Al, Fe, Cu, Zn and Mn, thus reducing the bad effects. Where the stability of complexes between humic acid-metal getting weaker in the order of $Al^{3+} > Fe^{3+} > Cu^{2+} > Mn^{2+} > Zn^{2+} \gg Mg^{2+} > Ca^{2+}$ (Tan, 2003). Nelvia (2009) reported that application of ameliorant Fe^{3+} and rock phosphates containing high Fe cation increased the stability of peat soil and reduced the carbon loss around 1.7 Mg of C $ha^{-1} year^{-1}$ (64%) in 5 cm of saturated condition, 1.3 Mg of C $ha^{-1} year^{-1}$ (58%) in two times of field capacity condition and 1.0 Mg of C $ha^{-1} year^{-1}$ (41%) in field capacity condition.

Dregs is the precipitate formed from liquid clarification process in the pulp mill recovery and no longer useful for the pulping process. Nelvia *et al.* (2008) reported that dregs contain polyvalent cations such as Al, Fe, Cu, Zn, Mn, Mo, and also contains other nutrients such as P, K, Ca, and Mg, the application of 15 tons dregs / ha of peatlands increases stover dry weight, dry weight of corn kernels per ear, 1000 grain weight seeds respectively 127%, 35% and 40% compared without dreg. This research aimed to study the potential of using dregs to reducing C-release (CO_2 and CH_4) and increasing growth and yield of rice on peat soil in unsaturation and saturation.

II. MATERIAL AND METHODS

This research was conducted from July to December 2010 in a greenhouse of Agriculture Faculty of Riau University. Soil chemical properties of peat soils materials and dregs was analyzed at soil laboratory of Soil Research Bogor. The peat soil material was taken at depths from 0 to 30 cm, with weathering rate saprik and dregs from Indah Kiat Pulp and Paper (IKPP) at Perawang, Riau.

The peat soil material was taken at depths from 0 to 30 cm, with weathering rate saprik at Kerumutan village, Pelalawan Regency, Riau Province, while dregs from from Riau Indah Kiat Pulp and Paper (IKPP) at Perawang, Riau, dregs chemical properties can be seen in Table 1. These experiments using split plot design, the main plot is the condition of the water (unsaturated and saturated) and the subplot is ameliorant dregs consisting of 4 levels (0, 10, 15 and 20 ton ha^{-1}), each combination was repeated 4 times.

Implementation of research: peat soil material equivalent of 2 kg dry weight oven 105 °C and dregs was mixed with appropriate treatment and then incubated at saturation and unsaturation condition for 1 month and then the IR-64 variety rice was planted. Basic fertilizers: Urea, TSP, and KCl each with a dose of 350, 150, and 150 kg ha^{-1} respectively.

To measure the flux of CO_2 and CH_4 , a chamber for trapping the gases made from the fiberglass with the size of 0.75 m x 0,20 m x 0,20 m, was used Syringes were used to take the samples of gases from the chamber. The samples were then put on the vacuum bottles. In this research, Gas Chromatography Shimadzu 14-B and Chromatopac Shimadzu C-R6A were used to determine the CO_2 and CH_4 emissions. The emissions were calculated by using following equation (Boer *et al.*, 1996):

$$\Phi_M = \{(\delta[CO_2/CH_4])\delta t \times h_U \times 16.123 (44.01) \times 273.2 \times (60/22.410)\}/(t_U + 273.2) \text{ mg m}^{-2} \text{ h}^{-1}$$

Where: $\delta[CO_2/CH_4]\delta t$ = change of the concentrations of CO_2 and CH_4 in chamber after the periode of t minute (s); h_U = the height of chamber; t_U = the average of air temperature in chamber; Value of 16.123 = the weight of CH_4 molecule, 44.01 = weight of CO_2 ; Value of 273.2 = temperature in Kelvin; 22.41 = volume of gas molecule; and Value of 60 means 60 minutes (1 hour). Other parameters were observed between: plant height 42 days after planting, the maximum tillers and productive tillers number, straw dry weight and grain dry milled weight.

III. RESULTS AND DISCUSSION

A. Chemical Composition of Peat

Several chemical characteristics of peats (Table 1) interesting to discuss. Although the total N is high, but the C/N ratio is very high, this means that N is a structural constituent of peat organic matter that is available N is low, thereby becoming a limiting factor for plant growth. cation exchange capacity (CEC) value is very high, but base saturation (BS) is very low, thus inhibiting the provision of nutrients, mainly K, Ca and Mg for plants.

The situation got worst because exchangeable Na, K, Ca and Mg is very low and therefore inhibit the growth and yield. Availability and total micro nutrient content are very low except for Fe is quite high, causing micro nutrient deficient for plants. According to Simbolon (2009) peat soil pH is very low, the availability of macro (N, P, K, Ca and Mg) nutrient are low and deficient micro (Cu, Zn, Mn, Fe, B and Mo) nutrient, CEC is very high but BS is low. Where the availability of Cu is the lowest compared to other micro nutrient because the Cu bounds to organic compounds functional groups such as carboxyl (COOH) and phenolic (-OH) to form organo—cation complex of Cu (chelate) that are not available for plants.

Several chemical characteristics (macro and micro nutrient) contained in dregs quickly available in peat, because hasil the extraction with 2% citric acid is almost equal to the extraction with mineral acids ($HClO_4$ and HNO_3 pa) (Table 2). Dissociation of H ions from organic compounds cause the concentration of H^+ ions on peat soil is very high, H^+ ions can hydrolyze dregs so that it dissolves quickly. Results of analysis of heavy metal content in the dregs (Pb, Cd, As, Hg, Co, Ni, Cr, Ag, Sn and Mo) total are very low (Table 3), are not including those identified B3 waste so it can be dumped in landfill light weight category.

TABLE I
CHEMICAL CHARACTERISTICS AND ASH CONTENT OF PEAT SOILS USED IN THIS RESEARCH

Chemical characteristics	Value	Chemical characteristics and ash content	Value
pH H ₂ O (1:5)	3.2	Base Saturation (%)	6
pH KCl (1:5)	3.0	Micro nutrient (Extract. DTPA)	
Organic-C (g kg ⁻¹)	437.3	Fe (µg/g)	475
Total-N (g kg ⁻¹)	6.5	Mn (µg/g)	1
C/N ratio	67.28	Cu (µg/g)	2
Exc.Ca (cmol (+)/kg)	2.27	Zn (µg/g)	2
Exc.Mg (cmol (+)/kg)	0.68	Micro nutrient (Extract. HNO ₃ + HClO ₄ pa)	
Exc.K (cmol (+)/kg)	0.22	Fe (µg/g)	3606
Exc.Na (cmol (+)/kg)	0.26	Mn (µg/g)	12.3
P ₂ O ₅ (µg/g) (Bray I)	135.4	Cu (µg/g)	3.1
P ₂ O ₅ (mg/100g) (HCl 25%)	32	Zn (µg/g)	4.8
CEC (cmol (+)/kg)	72.45	Ash content (%)	15.89

TABLE III
CHEMICAL CHARACTERIZATION AND MOISTURE OF DREGS

Chemical characteristics	Value	Chemical characteristics and ash content	Value
pH H ₂ O (1:5)	9.3	Macro nutrient (Extract. Citric Acid 2%)	1.8
Macro nutrient (Extract. HClO ₄ & HNO ₃ pa)		P ₂ O ₅ (g kg ⁻¹)	
P ₂ O ₅ (g kg ⁻¹)	2.0	K ₂ O (g kg ⁻¹)	3.1
K ₂ O (g kg ⁻¹)	3.1	CaO (g kg ⁻¹)	409.7
CaO (g kg ⁻¹)	410.3	MgO (g kg ⁻¹)	23.2
MgO (g kg ⁻¹)	23.9	Na (g kg ⁻¹)	25.9
Na (g kg ⁻¹)	26.8	S (g kg ⁻¹)	
S (g kg ⁻¹)	7.2	Micro nutrient (Extract. Citric Acid 2%)	6.4
Micro nutrient (Extract. HClO ₄ & HNO ₃ pa)		Fe (μg g ⁻¹)	3244
Fe (μg g ⁻¹)	5000	Mn (μg g ⁻¹)	914
Mn (μg g ⁻¹)	989	Cu (μg g ⁻¹)	105
Cu (μg g ⁻¹)	127	Zn (μg g ⁻¹)	206
Zn (μg g ⁻¹)	224	moisture (%)	15

TABLE IIIII
HEAVY METAL CONTENT OF DREGS

Chemical characteristics	Value	Chemical characteristics	Value
Extraction (HClO ₄ + HNO ₃ pa)		Extraction (citric acid 2%)	
Pb (μg g ⁻¹)	8.9	Pb (μg g ⁻¹)	0.1
Cd (μg g ⁻¹)	0.2	Cd (μg g ⁻¹)	nm
As (μg g ⁻¹)	3.8	As (μg g ⁻¹)	nm
Hg (μg g ⁻¹)	0.23	Hg (μg g ⁻¹)	nm
Co (μg g ⁻¹)	1.7	Co (μg g ⁻¹)	1.5
Ni (μg g ⁻¹)	98.6	Ni (μg g ⁻¹)	98.5
Cr (μg g ⁻¹)	167	Cr (μg g ⁻¹)	120
Se (μg g ⁻¹)	355	Se (μg g ⁻¹)	169
Ag (μg g ⁻¹)	nm	Ag (μg g ⁻¹)	nm
Sn (μg g ⁻¹)	nm	Sn (μg g ⁻¹)	nm
Mo (μg g ⁻¹)	nm	Mo (μg g ⁻¹)	nm

B. The effect of application of ameliorant dregs in the peat at saturation and unsaturation condition on growth and yield of rice

The application of dregs improves plant growth (plant height, number of maximum tillers, number of productive tillers and weight of dry straw) and increase the yield of rice (weight of dry milled grain) compared without dreg both at unsaturated condition and saturated conditions (Table 4 and Fig. 1). Rice can not grow well in peat, which is not applied ameliorant dreg, the cause is the low nutrient availability and high content of phenolic acids that are toxic to plants. Tim Sintesis Kebijakan (2008) reports that rice plants grown in peat soil with a thickness of over 2 m deficient Cu failed to form a grain. The peat composition is dominated by lignin of 65% to 80% and 78% to 93% for the peats of Jambi and Central Kalimantan respectively (Sabiham, 2010), and 71,46% for the peats of Riau (Nelvia, 2009). Orlov (1995) showed the processes of lignin disintegration that result in several derivate phenolic acids. Phenolic acids are more phytotoxic for plants and causes stunted plant growth (Tsutuki, 1984; Stevens, *et al.*, 1994; Dohong and Sabiham, 2001), influence the biochemical and physiological processes of plants and nutrients uptake by plant influence the biochemical and physiological processes of plants and nutrients uptake by plant (Driessen, 1978). Tsutuki *et al.*

(1994) stated that the concentration of phenolic acids at the range of 0.6 to 3.0 mM could hamper the root growth of rice up to 50%. Todano *et al.* (1992) reported that derivate phenolic acids, such as ferulic, synapic, p-cumaric, and p-hydroxybenzoic acids are phytotoxic for rice, particularly during the first stage of plant growth. He also mentioned that ferulic acid in peat is more toxic compared to the other derivate phenolic acids.


TABLE IVV
THE EFFECT OF AMELIORANT DREGS IN THE PEAT ON THE PLANT HEIGHT, MAXIMUM AND PRODUCTIVE TILLERS NUMBER AND WEIGHT OF DRY STRAW AND MILLED DRY GRAIN OF RICE

Water condition	Dregs (ton ha ⁻¹)	Plant height (cm)	maximm tillers (number pot ⁻¹)	productive tillers (number pot ⁻¹)	dry straw (g pot ⁻¹)	milled dry grain (g pot ⁻¹)
Saturated	0	47 c	20 d	17 c	20 c	20.0c
	10	66 a	33 c	23 bc	42 b	35.4a
	15	67 a	45 ab	31 a	61 a	54.8b
	20	67 a	47 a	30 a	62 a	54.9b
Un-saturated	0	52 b	20 d	19 c	27 c	29.8 c
	10	64 a	36 abc	31 a	43 b	53.7 b
	15	68 a	35 bc	28 ab	46 b	53.0 b
	20	66 a	36 abc	28 ab	52 b	45.3 a

Note: The numbers in the same columns which followed the same lowercase letter are not significantly different at a 5% DNMR test.



a. Unsaturation condition



b. Saturation condition

Fig. 1. The growth of rice at vegetative and generative phase in unsaturated (a) and saturated condition (b)

The application dreg 10 t ha⁻¹ increase the maximum number of tillers, number of productive tillers, straw dry weight and the weight of milled rice by 65, 35.29, 110 and 75%, respectively compared without dreg on saturated condition, whereas the increase is greater unsaturated conditions about 80, 63, 59 and 80%, respectively (Table 4). This is due to the improvement of the condition of the peat soil chemistry by ameliorant dreg in the form increase nutrient availability and decreasing the solubility of phenolic acids. Its caused by the dregs contain macro (P, K, Ca, Mg, S) and micro (Fe, Cu, Zn, Mn, Mo) nutrients or polivalen cations and quickly available in peat (Table 2). Reduced

solubility of phenolic acids occurs due to the formation of complex compounds of phenolic acids with cations Fe, Cu, Zn, Mn, Ca and Mg are dissolved from the dreg. Sabiham (2010) reported that the concentration of derivate phenolic acids namely: ferulic, syanapic, p-cumaric, vanilic, syringic and p-hydroxybenzoic acids in peats decreased with the addition of mineral soil or basic slag, or the ombination of both materials.

C. The effect of application of ameliorant dregs in the peat soil at unsaturation and saturation condition on carbon emission

The Cabon-release in the forms of CO₂ and CH₄ fluxes in saturated conditions is smaller than unsaturated conditions, the application of dregs 10 t ha⁻¹ in saturated condition can reduce CO₂ emissions about 18.19% and CH₄ emissions about 93.71% compared without dreg, otherwise the application dreg 10, 15 and 20 t ha⁻¹ in unsaturated conditions increase the production of CO₂ and CH₄ (Table 5). Unsaturated conditions (aerobic) conditions where the peat is available high due to the high diffusion, so the more active microorganisms both the type and amount of the anaerobic conditions, the result would accelerate the process of decomposition of organic matter. While in saturated conditions (anaerobic) only anaerobic bacteria can live, thus CO₂ as a result of respiration and decomposition of organic matter will be higher in unsaturated than saturated conditions. Boer *et al.* (1996) reported that the amount of CH₄ emission rate depends on soil water conditions, flooded peatlands emit CH₄ greater than the land is not flooded. Research by Sabiham and Sulistyono (2000) in the laboratory showed that the highest CO₂ production obtained in aerobic incubation and significantly different with anaerobic incubation, while the highest CH₄ production was obtained on anaerobic incubation and significantly different with aerobic incubation.

TABLE V

THE EFFECT OF APPLICATION OF AMELIORANT DREGS IN THE PEAT SOIL AT SATURATION AND UNSATURATION CONDITION ON THE CO₂ AND CH₄ PRODUCTION

Water condition	Dregs (ton ha ⁻¹)	CO ₂ production (mg pot ⁻¹ h ⁻¹)	CH ₄ production (mg pot ⁻¹ h ⁻¹)
saturation	0	8.12 a	29.70 a
	10	6.64 b	1.87 c
	15	8.38 a	3.07 bc
	20	7.31 b	2.68 bc
unsaturation	0	18.72 b	6.69 b
	10	29.71 b	7.61 b
	15	52.57 a	19.39 a
	20	15.64 b	5.78 b

Note: The numbers in the same columns which followed the same lowercase letter are not significantly different at a 5% DNMRT test

The decrease in emissions of CO₂ and CH₄ in the saturated condition by administering dreg 10 t ha⁻¹ is because dreg containing polyvalent cations and release such as Fe and Cu to the soil (Table 2). Fe and Cu cations form complex compounds between organic compounds with Fe or

Cu cations, that is stable so it can not be decomposed by microbes. Giving dreg with the higher dose (15-20) has stimulate more rapid decomposition of organic matter, especially in unsaturated conditions, while the formation of complex organic compounds-metal running over time, The result is an increase in emissions of CO₂ and CH₄. Research Sabiham and Sulistyono (2000) in the laboratory showed that administration of Fe³⁺ cations as much as 5% maximum erapan can reduce 22.94% 23.01% CO₂ and CH₄ in the peat soil of the area Dendang Jambi and 27.67% and 32.97% CO₂ CH₄ in the peat soil of the Sampit, Central Kalimantan. The addition of 15 and 30 g of Fe(OH)₃ per kg of soil to lower the total CH₄ emissions by 43% and 84% during the growth of rice (Jackel and Schnell 2000). Where the stability of complexes between humic acid-metal getting weaker in the order of Al³⁺ > Fe³⁺ > Cu²⁺ > Mn²⁺ > Zn²⁺ >> Mg²⁺ > Ca²⁺ (Tan, 2003).

IV. CONCLUSIONS

The application of dregs improves plant growth and increase the yield of rice compared without dreg both at unsaturated condition and saturated conditions. The application dreg 10 t ha⁻¹ increase the maximum number of tillers, number of productive tillers, straw dry weight and the weight of milled rice by 65, 35, 110 and 75%, respectively compared without dreg on saturated condition, whereas the increase is greater unsaturated conditions about 80, 63, 59 and 80%, respectively.

The Cabon-release in the forms of CO₂ and CH₄ fluxes in saturated conditions is smaller than unsaturated conditions, the application of dregs 10 t ha⁻¹ in saturated condition can reduce CO₂ and CH₄ emissions about 18.19% and 93.71% compared without dreg, otherwise the application dreg 10-20 t ha⁻¹ increase the production of CO₂ and CH₄ in unsaturated conditions.

REFERENCES

- [1] Boer, R., I. Nasution, I. Las and A. Bey. 1996. Emisi metan dari lahan gambut sejuta hektar, Kalimantan Tengah (Methane emission from the one-million-hectare peatlands, Central Kalimantan). *J. Agromet* 12 (1&2) 1996/1997. Pp. 31-38.
- [2] Dohong, S. S. Sabiham, 2001. Several derivate of phenolic acids on Central Kalimantan on different environment of peat formation. *Agrista*. Vol 5 (3): 197-203.
- [3] Driessen, P.M. 1978. Peat soils. In: *IRRI. Soil and rice*. IRRI. Los Banos. Philippines. Pp: 763-779.
- [4] Hooijer, A, Page, S., Canadell, J., Silvius, M. Kwidjkk, J. Wosteten, H. and Jauhiainen J. 2010. Curren and future CO₂ emission from drained peatlands in Southeast Asia. *Biogeosciences*, 7, 1505-1514
- [5] Jackel, U and S. Schnell. 2000. Suppression of methane emission from rice paddies by ferric iron fertilization. *Soil Biology & Biochemistry*. 32:1811-1814.
- [6] Nelvia, Rosmimi, Rini and R. Frizdew. 2008. Peningkatan produktivitas lahan gambut dengan pemberian amelioran dregs (Limbah bagian Recauticizing pabrik pulp) berkadar kation polivalen tinggi. Dalam *Prsiding Semiloka Nasional*. Departemen Ilmu Tanah dan Sumberdaya Lahan IPB. Bogor. 22-23 Desember 2008. Pp 497-506.
- [7] ----- . 2009. Kandungan fosfor tanaman padi dan emisi karbon tanah gambut yang diaplikasi dengan amelioran Fe³⁺ dan fosfat alam pada beberapa tingkat pemberian air. *Jurnal Tanah tropika* Vol. 14 (3): 195-204.
- [8] Orlov, D.S. 1995. Humic substanscess of soils and general theory of humification. *A.A. Balkema Publ. USA*.

- [9] Sabiham, S. and N.B.E. Sulistyono. 2000. Kajian beberapa sifat inheren dan perilaku gambut: Kehilangan karbondioksida (CO₂) dan metana (CH₄) melalui proses reduksi-oksidasi. *J. Tanah Tropika*. V (10): 127-135.
- [10] _____. 2010. Properties of Indonesian peat in relation to the chemistry of carbon emission. In: *Proceeding of International Workshop on Evaluation and Sustainable Management of Soil Carbon Sequestration in Asian Countries*. Pp: 205-216
- [11] Simbolon, H. 2009. Peat swamp forest ecosystem: An important ecosystem on regional land use planning. In *Scientific Exploration and Sustainable Management of Peat Land Resources in Giam Siak Kecil-Bukit Batu Biosphere reserve*. Riau. Pp: 165-174.
- [12] Stevens, D.P., M.J. McLaughlin and A.M. Alston. 1994. Are Aluminum-Fluoride complexes Phytotoxic. 15th World Congress of Soil Science. Acapulco. Mexico
- [13] Tadano, K. Yonebayashi and N.Saito 1992. Effect of phenolic acids on the growth and occurrence of sterility in crop plants. Pp:358-369. In: K.Kyuma, P.Vajarnsorn and A.Zakaria (eds) *Coastal lowland ecosystems in southern Thailand and Malaysia*. Showado-printing co. Skyoku-Kyoto
- [14] Tan, K.H. 2003. *Humic Matter in the soil and the environment; Principles and Controversies*. Marcel Dekker, Inc. new York. USA. P 359.
- [15] Tim Sintesis Kebijakan. 2008. Pemanfaatan dan konservasi ekosistem lahan rawa gambut di Kalimantan. *Pengembangan Inovasi Pertanian* 1 (2): 149-156.
- [16] Tsutsuki, K. 1984. Volatile products and low molecular-weight products of the anaerobic decomposition of organic matter. pp :329-343. *International Rice Research Institute, Soil Organic Matter*.
- [17] Wahyunto, Ritung S., Suparto and Subagjo H. 2005. Peat land Distribution and Carbon Content in Sumatera and Kalimantan. *Wetland International- Indonesia Program and Wildlife Habitat Canada (WHC)*. Bogor-Indonesia
- [18] Yagi, K and K. Minami. 1990. Effect of organic matter application on methane emission from some Japanese paddy field. *Soil Sci. Plant Nutr.* 36(4):599-610.